

# Adult Pneumococcal Vaccine Timing


## Age 65 Years and Older

\*If PCV13 was given before age 65, no additional PCV13 is needed

No history of  
pneumococcal  
vaccine



1 Year (8 Weeks for groups B & C  
as defined below)




Received  
PPSV23  
before  
age 65



1 Year (8 Weeks for groups B & C  
as defined below) and 5 years  
after prior dose of PPSV23



Received  
PPSV23  
at age 65  
or older



## Age 19-64 Years With Underlying Medical Condition(s)

\*Prior doses count towards doses recommended below and do not need to be repeated.

\*If PPSV23 was given previously: (1) Wait one year before giving PCV13 (2) For group B, wait at least five years before giving a second dose of PPSV23. \*No more than two doses of PPSV23 recommended before 65th birthday and one dose thereafter.


### A. Smoker, long-term facility resident, or chronic conditions:

- Heart Disease (excluding hypertension)
- Lung Disease (including asthma)
- Liver Disease (including cirrhosis)
- Diabetes • Alcoholism





### B. Immunocompromised

- HIV Infection
- Chronic Renal Failure
- Nephrotic Syndrome
- Asplenia



### C. CSF Leaks or Cochlear Implants



For further information,  
see: <http://www.cdc.gov/vaccines/vpd-vac/pneumo/default.htm>